DH CIRCULAR No. 44/2017

DH 827/2017

21 June 2017

Attention All: Consultants

Medical Officers Pharmacists

Pharmacy Technicians

Nurses

Re: Mitomycin C 40mg Vials

It is being brought to your attention that mitomycin C 40mg vials are now available on the Government Formulary List for in-patient use.

Mitomycin C 40mg vials are presented as powder for solution for injection to be used for bladder instillation and to be administered intravesically via Electromotive Drug Administration (EMDA).

Mitomycin C 40mg vials are protocol regulated by protocol 138 (Annex I) and can be prescribed by Consultant Urologists for superficial bladder tumours as per said protocol.

For your attention please.

Dr. Denis Vella Baldacchino Chief Medical Officer

138

Mitomycin C 40mg Vials

(To be administered intravesically via Electromotive Drug Administration [EMDA])

Prescriber criteria: Consultant Urologist

In-patient use:

1. Neoadjuvant use prior to resection of superficial bladder tumours

To be used as a perioperative single dose instillation before Transurethral Resection of Bladder Tumour (TURBT)

2. Management of high risk superficial bladder tumours

To be used either:

- in combination with BCG during the induction and maintenance course OR
- on its own during the induction and maintenance course when BCG is not available (as a stop gap measure), or in patients who are either hypersensitive to BCG or have a past history of TB